

Table 2S. Influence of P application rate on shoot dry mass [g per plant] at tillering stage and reproductive stage of selected rice varieties. Each value represents mean of 3 replications.

Variety	Shoot dry mass [g per plant]											
	Tillering stage						Reproductive stage					
	P application rate (kg ha ⁻¹)											
	0	15	30	45	60	Mean	0	15	30	45	60	Mean
Akhanphou	0.18	6.15	14.57	23.35	34.45	15.74	-	22.74	38.10	63.61	79.90	51.08
MTU 1010	0.16	3.65	17.57	25.99	33.86	16.24	-	21.62	31.11	44.47	61.93	39.78
RP BIO 226	0.12	0.22	12.47	26.66	34.81	14.86	-	11.10	22.45	41.42	56.49	32.87
Swarna	0.18	11.29	20.48	42.19	53.96	25.62	-	26.55	40.25	61.95	77.46	51.55
Mean	0.16	5.33	16.27	29.55	39.27		-	20.50	32.98	52.86	68.94	
LSD(Treatment)			1.72	(P<0.05)					1.59	(P<0.05)		
LSD (Varieties)			1.53	(P<0.05)					1.42	(P<0.05)		
LSD (T X V)			3.43	(P<0.05)					3.17	(P<0.05)		