

Table 3S. Influence of P application rate on root dry mass [g per plant] at tillering stage and reproductive stage of selected rice varieties. Each value represents mean of 3 replications.

Variety	Root dry mass [g per plant]											
	Tillering stage						Reproductive stage					
	P application rate (kg ha ⁻¹)											
	0	15	30	45	60	Mean	0	15	30	45	60	Mean
Akhanphou	0.03	0.48	0.62	1.41	1.81	0.87	-	0.93	7.25	10.97	16.24	8.84
MTU 1010	0.03	0.28	0.50	0.94	1.94	0.74	-	0.88	7.39	10.75	14.63	8.41
RP BIO 226	0.03	0.04	0.32	0.79	1.73	0.58	-	0.22	6.02	9.97	12.66	7.21
Swarna	0.04	0.30	0.87	1.58	2.36	1.03	-	3.21	8.40	15.20	12.91	9.93
Mean	0.03	0.27	0.57	1.18	1.96		-	1.31	7.26	11.72	14.11	
LSD(Treatment)			0.13	(P<0.05)					0.88	(P<0.05)		
LSD (Varieties)			0.11	(P<0.05)					0.79	(P<0.05)		
LSD (T X V)			0.25	(P<0.05)					1.76	(P<0.05)		