

Table 4S: Influence of P application rate on leaf and root P concentration [mg g⁻¹ DM] at tillering stage of selected rice varieties. Each value represents mean of 3 replications.

Variety	Leaf P concentration [mg g ⁻¹ DM]						Root P concentration [mg g ⁻¹ DM]					
	P application rate (kg ha ⁻¹)											
	0	15	30	45	60	Mean	0	15	30	45	60	Mean
Akhanphou	0.59	1.44	1.34	3.53	3.43	2.07	0.39	0.67	1.05	1.56	1.74	1.08
MTU 1010	0.26	1.10	1.28	3.81	3.13	1.92	0.31	1.08	1.06	0.94	2.06	1.09
RP BIO 226	0.48	2.08	1.35	3.67	4.14	2.34	0.29	1.01	0.99	1.49	1.94	1.14
Swarna	0.54	1.18	2.81	4.26	3.44	2.44	0.32	1.66	1.48	1.49	2.54	1.50
Mean	0.47	1.45	1.69	3.82	3.53	2.19	0.33	1.11	1.14	1.37	2.07	1.20
LSD(Treatment)			0.11	(P<0.05)						0.11	(P<0.05)	
LSD (Varieties)			0.10	(P<0.05)						0.10	(P<0.05)	
LSD (T X V)			0.23	(P<0.05)						0.23	(P<0.05)	