



## Gas exchange and time to reach maximum rate of photosynthetic rate and their relationship with whole-plant traits in sugarcane in water abundant Louisiana, USA

P.Z. ELLSWORTH<sup>+</sup> , P.M. WHITE JR. , and J. TODD 

USDA-ARS Sugarcane Research Unit, 70360 Houma, LA, USA

### Abstract

Variety development of sugarcane (*Saccharum* spp. hybrids) is necessary to continue improving sugar yields and selecting photosynthetic traits can improve sugar production through increased carbon inputs. In this study, gas exchange and whole-plant measurements were made on 55 sugarcane genotypes in Louisiana. Variation in the relationship between photosynthetic rate and stomatal conductance suggests that sugarcane exhibits variation in both photosynthetic capacity and CO<sub>2</sub> substrate availability. Genotypes that reached maximum photosynthetic rate (TRMPR) in the gas-exchange cuvette more quickly had greater CO<sub>2</sub> assimilation during transitory periods. Temporary shading and fluctuating light are common transitory conditions in the field, so increasing TRMPR can improve photosynthesis in water-abundant regions. Canopy leaf area was positively correlated with stalk mass, but gas-exchange traits were not correlated with whole-plant traits. A better understanding of the relationship between leaf and whole-plant traits is necessary to identify physiological traits that lead to increased genetic gain.

**Keywords:** gas exchange; phenotyping; photosynthesis; stomatal conductance; stomatal propensity to remain open; sugarcane.

### Introduction

Sugarcane (*Saccharum* spp. hybrids) produces nearly 80% of sugar globally, is a major source of ethanol, and is the largest crop by biomass in the world (ISO 2023). In Louisiana, it is the second largest agricultural crop covering 251,000 ha, and valued at over \$3.9 billion to the Louisiana economy in 2020 (ASCL 2023). The current efforts of the USDA-ARS Sugarcane Research Unit, Louisiana State University AgCenter, and the American Sugar Cane League are focused on producing new sugarcane cultivars that improve aboveground biomass, early season sugar accumulation, total recoverable sugar, disease resistance, and resistance to sugarcane borers (Hale *et al.* 2022). These efforts have increased sugar recovery from 5.8 to 11.7% from 1890 to 2010 and

increased average tons of sugar per hectare from 2.5 to 8.8 over the same time period (Hale *et al.* 2022). Further progress in sugarcane variety development can be made by discovering additional physiological traits and high throughput phenotyping methods. Phenotyping of traits, especially physiological traits, requires, first, identifying specific traits that have a clearly defined role in improving sugarcane, are heritable, and exhibit genetic variation within the breeding population (Araus *et al.* 2014). Second, a preferably high throughput phenotyping method needs to be developed that is accurate, repeatable, and rapid. Selection of simple traits improve selection efficiency because it improves the relationship between genetic gain and phenotyping where genetic gain is the increase in yield or performance over time through the efforts of artificial selection and is a function of selection intensity,

### Highlights

- Substantial variation in gas-exchange traits in sugarcane
- Short time to reach maximum rate of photosynthetic rate increases photosynthesis
- Canopy leaf area and stalk mass positively correlated

Received 20 September 2023

Accepted 5 March 2024

Published online 20 March 2024

<sup>+</sup>Corresponding author

e-mail: patrick.ellsworth@usda.gov

**Abbreviations:** C<sub>a</sub> – ambient CO<sub>2</sub> concentration; C<sub>i</sub> – intercellular CO<sub>2</sub> concentration; E – transpiration rate; g<sub>s</sub> – stomatal conductance; P<sub>N</sub> – net photosynthetic rate; TMPR – time to reach maximum photosynthetic rate.

**Conflict of interest:** The authors declare that they have no conflict of interest.

selection accuracy, and genetic variance relative to the years per breeding cycle (Araus *et al.* 2018). For example, greater genetic gains can be made through identifying photosynthetic traits that have high phenotypic variation and are under genetic control because of the potential of increasing the carbon available for sucrose production.

Photosynthesis has long captured the attention of plant physiologists and breeders alike as a source of improved plant productivity (von Caemmerer and Furbank 2003, Long *et al.* 2006, von Caemmerer *et al.* 2009, Evans 2013, Ort *et al.* 2015, Lawson *et al.* 2018). Most progress in increasing yield has focused on increasing carbon allocation to the harvested portion such as increasing early sucrose production in Louisiana and increasing aboveground biomass (harvest index) (Araus *et al.* 2014, 2018). Harvest index and carbon allocation to sucrose will reach their maximum threshold where the infrastructural needs of leaves, roots, and stems cannot be reduced further without negative feedback effects on sugar production. In contrast, increasing carbon inputs through improved photosynthesis has not undergone direct selection and remains a potentially major source of improvement to sugarcane growth and sugar production. To select for photosynthetic traits in Louisiana sugarcane such as leaf gas-exchange traits, phenotypic variability across genotypes must be present. Leaf gas exchange characterizes the movement of CO<sub>2</sub> and H<sub>2</sub>O through leaf stomata where CO<sub>2</sub> diffuses into the leaf for photosynthetic assimilation and H<sub>2</sub>O vapor exits the leaf. Gas-exchange measurements quantify CO<sub>2</sub> assimilation in photosynthesis and transpiration water loss and stomatal conductance at the leaf level and provide insight into leaf behavior that influences photosynthesis (Long and Bernacchi 2003). For example, gas exchange can shed light on stomatal responses to various biochemical and environmental controls and the environmental and physiological factors that influence components of photosynthesis such as enzyme concentrations and activity (von Caemmerer 2000). Because these traits represent the only influx of carbon through photosynthesis and the major source of water loss in the plant, there is an inherent relationship between leaf-level photosynthetic traits and whole-plant growth and water-loss traits.

Relationships between leaf and whole-plant traits provide insight into how leaf-level photosynthesis and transpiration influence whole-plant behavior. Leaf-level photosynthesis is the source of all carbon assimilated into plant tissue and the source of most water loss through transpiration, but it is also how carbon is used and retained in the plant that determines crop productivity and yields. Therefore, it is necessary to understand both basic photosynthesis and transpiration as leaf traits and their interaction with whole-plant traits such as canopy area and stalk traits that interconnect photosynthesis and transpiration with yield and productivity (Farquhar *et al.* 1989). However, the complex nature of whole-plant traits often obscures the role of leaf-level gas exchange in whole-plant traits, especially in perennial and woody species, which makes direct connections between leaf and whole-plant scales difficult (Medrano *et al.* 2015). Nonetheless, multiple studies have found relationships

between gas-exchange traits and growth and water use at the whole-plant level (Ellsworth *et al.* 2017, 2020; Feldman *et al.* 2018, Leakey *et al.* 2019).

In this study, 55 sugarcane genotypes were measured for various leaf-level traits including net photosynthetic rate ( $P_N$ ) and transpiration ( $E$ ), stomatal conductance ( $g_s$ ), leaf area, and time to reach maximum photosynthetic rate (TRMPR) after being placed in the gas-exchange cuvette, and whole-plant traits such as cane biomass volume and canopy leaf area. We hypothesized that because of the large genetic variation that is found in sugarcane genotypes, these leaf-level traits would have substantial phenotypic variation across genotypes and that significant relationships would be present between leaf and whole-plant level photosynthetic and water loss traits. The objectives were to (1) measure the phenotypic variation of these traits and to determine if there is a significant genotypic effect, (2) develop relationships between gas exchange and whole-plant traits and compare gas exchange and whole-plant traits between commercial cultivars and genotypes in the breeding program.

## Materials and methods

**Study site and sugarcane genotypes:** The plants used in this study were grown in the field on the USDA-ARS Ardoyne farm in Louisiana, USA. All sugarcane plants were in plant cane crop and were grown in the same field on 1.8 m-spaced rows in a Cancienne silty clay loam soil under rainfed conditions. Plots were 2 adjacent, 7.6-m long row sections. A total of 55 sugarcane genotypes were used in this study (listed in Table 1S, *supplement*). The genotypes were either commercially released cultivars (16) or mid-stage genotypes in the breeding program (39) whose parents were a combination of commercially released cultivars and late-stage breeding genotypes from the USDA-ARS variety development program. These genotypes were chosen because they represented the major commercial genotypes and an entire cohort of new genotypes, some of which may be released as a commercial cultivar.

**Gas exchange:** Stalks from the field plots were cut at the base and immediately placed in a bucket of water and taken to the lab. Once in the lab, the cut end of the stalk was re-cut while submerged in water. Gas-exchange measurements were made with *LI-6800* (*LI-COR*, Lincoln, NE, USA). For gas-exchange measurements, the youngest, fully expanded leaf was placed in a 3 cm by 3 cm gas-exchange cuvette (*6800-12A*). The conditions in the cuvette were the following: 1,500  $\mu\text{mol}(\text{photon})\text{ m}^{-2}\text{ s}^{-1}$  of PAR (90% red and 10% blue light), 30°C leaf temperature, VPD of  $1.12 \pm 0.16$  kPa, flow was set at  $600\ \mu\text{mol s}^{-1}$ , mean relative humidity of  $74.3 \pm 3.6\%$ , and sample CO<sub>2</sub> concentration of  $400\ \mu\text{mol mol}^{-1}$ . The midsection of the leaf including the midvein was placed in the cuvette. Measurements were logged every minute during the entire time that the leaf was in the cuvette, and the sample and reference infrared gas analyzers were matched every 10 min. The leaf was removed once  $P_N$  and  $g_s$  remained constant for at least 4 min.

To verify that measuring gas exchange on leaves from cut stalks did not affect the measurements, gas-exchange measurements were made on 25 potted sugarcane plants from a total of nine genotypes. Leaves were inserted in the gas-exchange cuvette and placed under the same conditions stated above. Once  $P_N$  and  $g_s$  remained constant for 4 min, the leaves were removed from the cuvette. At this point the stalk was cut, and the cut end was placed in a bucket of water then recut under water. The same leaf but in a different location was placed in the gas-exchange cuvette again and remeasured under the same conditions. These paired measurements were compared to determine if  $P_N$ ,  $E$ , and  $g_s$  changed after cutting the stalk and more importantly if this change varied with genotype. In a two-way repeated measures ANOVA with factors genotype and treatment (uncut vs. cut stalk), genotype was significant for  $P_N$ ,  $E$ , and  $g_s$ , and the interaction was not significant ( $P > 0.05$ ). From these results, we conclude that cutting the stalk did not affect gas-exchange measurements and especially did not affect genotypes differentially or systematically. The model II regression slope of the line between gas exchange values of leaves from uncut (x-axis) and cut stalks (y-axis) was 1.1 for  $P_N$ , 0.94 for  $g_s$ , and 0.83 for  $E$ , showing that the gas-exchange values were similar before and after cutting the stalk. For time to reach maximum net photosynthetic rate (TRMPR) in the two-way repeated measures ANOVA, interaction was not significant ( $P > 0.1$ ), meaning that the response in TRMPR was not dependent on genotype. Therefore, measuring gas exchange of leaves from cut stalks was considered appropriate.

TRMPR was measured as the time when the leaf was placed in the gas-exchange cuvette until the leaf reached the maximum  $P_N$  and remained constant for 4 min without increasing further. Cumulative CO<sub>2</sub> assimilation [ $\mu\text{mol}(\text{CO}_2) \text{ m}^{-2}$ ] and transpiration [ $\text{mmol}(\text{H}_2\text{O}) \text{ m}^{-2}$ ] represented the total CO<sub>2</sub> assimilation and water loss per m<sup>2</sup> during the time the leaf was reaching maximum photosynthesis. Each minute log was assumed to represent the entire minute, so the sum of one-minute logs of  $P_N$  and  $E$  from the time that the leaf was placed in the cuvette until it reached its maximum  $P_N$  was multiplied by 60 to calculate them over the entire minute time frame.

**Whole-plant measurements:** Stalks were harvested by cutting them flush with the soil surface. The cut ends of the stalks were immediately placed in a bucket of water and stored in water until they were processed. The photosynthetically active, non-senesced leaves were removed from each stalk, and leaf area was measured using a portable leaf area meter, *LI-COR 3000C* (*LI-COR*, Lincoln, NE, USA). Then the leaves were placed in a paper bag and weighed in aggregate for the fresh mass and put in a drying oven at 60°C until their mass did not change further. At that point, the dry mass was recorded. The number of nodes were counted on the stalk, and total length and diameter at the basal end and middle of the stalk were measured. The stalk was weighed prior to being placed in a drying oven at 60°C and again once the mass was no longer decreasing. Cane volume was

calculated as the volume of a cylinder using the mean cane diameter and cane height.

**Statistical analysis:** All statistics were done in *R* (*R Core Team 2021*). Genotype effect was determined by one-way ANOVA of each trait. Differences between commercial cultivars and mid-stage genotypes in the breeding program were tested using *t*-tests of the genotype means. Type II linear regressions were used to calculate relationships between leaf and whole-plant traits. Using *lmodel2* (v. 1.7-2) package, model II regression analysis (standard major axis) was used as a more appropriate regression analysis than ordinary least squares regression for all linear regressions because neither independent nor dependent variables were controlled, both varied naturally with their own associated error, and the physical units of both variables were not the same.

## Results

**Gas-exchange traits:** Gas-exchange measurements [ $P_N$ ,  $g_s$ ,  $E$ , ratio of intercellular to ambient CO<sub>2</sub> concentrations ( $C_i/C_a$ ),  $C_i$ ] were consistent across replicates in each genotype in that standard errors were only 4–6% of the measured values, showing that these traits were reliable at the genotype level. All traits were significantly different across genotypes, showing a significant genotype effect (Table 1, Fig. 1). Variation across sugarcane genotypes was relatively high in gas-exchange traits, in that maximum values were 1.37 to 1.78 times that of the minimum values (Table 1). The commercial cultivars and unreleased genotypes in the breeding program did not differ in  $P_N$  and  $E$ , but slight differences were found in  $C_i/C_a$  and  $g_s$ , where commercial cultivars were slightly higher in both ( $P < 0.01$  and 0.05, respectively; Table 1). The mean maximum  $P_N$  reached in each individual leaf was  $30.5 \pm 2.2$  (SD)  $\mu\text{mol}(\text{CO}_2) \text{ m}^{-2} \text{ s}^{-1}$  with 38.9 (L 17-428) being the highest value measured.  $E$  and  $g_s$  were rather high for a C<sub>4</sub> grass, reaching  $3.51 \text{ mmol}(\text{H}_2\text{O}) \text{ m}^{-2} \text{ s}^{-1}$  and  $0.356 \text{ mol}(\text{H}_2\text{O}) \text{ m}^{-2} \text{ s}^{-1}$  (L 17-428), respectively, and averaging  $2.68 \pm 0.26 \text{ mmol}(\text{H}_2\text{O}) \text{ m}^{-2} \text{ s}^{-1}$  and  $0.253 \pm 0.029 \text{ mol}(\text{H}_2\text{O}) \text{ m}^{-2} \text{ s}^{-1}$ , respectively.

Across genotypes mean  $P_N$  was correlated with mean  $g_s$  as expected, suggesting an increase in  $P_N$  by increasing CO<sub>2</sub> substrate availability, but there was considerable variation across genotypes in this relationship in that  $R^2$  was only 0.51 (Fig. 2A). While most genotypes followed the expected positive relationship, a few genotypes fell off the regression line considerably. L 01-428, Ho 17-738, and HoCP 18-878 had high  $P_N$  and low  $g_s$ , suggesting higher photosynthetic capacity instead of increased CO<sub>2</sub> substrate availability, and HoCP 20-527, HoCP 20-548, and L 14-267 had low  $P_N$  and high  $g_s$ , suggesting lower photosynthetic capacity. Nearly all genotypes followed a general positive relationship between  $g_s$  and  $C_i/C_a$  followed a similar relationship ( $C_i/C_a = 1.034 g_s + 0.215$ ,  $P < 0.0001$ ,  $R^2 = 0.61$ ). L 01-428, Ho 17-738, and HoCP 18-878 had 7–12% lower  $C_i/C_a$  than expected based on  $g_s$ , further indicating increased photosynthetic capacity, and HoCP 20-527, HoCP 20-548, and L 14-267 had 7–11% lower  $C_i/C_a$  than expected, further indicating lower

Table 1. *ANOVA* and *t*-test table. 55 genotypes were measured including 16 commercial cultivars and 39 unreleased genotypes. Differences across genotypes were calculated using one-way *ANOVAs*. *t*-tests were used to determine if commercially released cultivars and mid-stage genotypes in the USDA-ARS breeding program were significantly different. Min. and Max. represent minimum and maximum values measured in the study. Canopy  $P_N$ ,  $E$ , and  $g_s$  represent these traits scaled to the entire canopy leaf area. \* $g_s$  was greater in commercial cultivars (0.266) than unreleased genotypes (0.249). † $C_i/C_a$  was greater in commercial cultivars (0.50) than unreleased genotypes (0.47). ‡ $C_i$  was greater in commercial cultivars (198.8) than unreleased genotypes (188.2).

Factor	Min.	Max.	Mean $\pm$ SE	One-way <i>ANOVA</i> (genotype)		<i>t</i> -test (type)	
				$F_{\text{ndf,ddf}}$	<i>P</i>	$T_{\text{df}}$	<i>P</i>
$P_N$ [ $\mu\text{mol m}^{-2} \text{s}^{-1}$ ]	26.23	38.90	30.46 $\pm$ 0.29	3.253 <sub>54,287</sub>	< 0.0001	1.966 <sub>1,287</sub>	0.16
Canopy $P_N$ [ $\mu\text{mol s}^{-1}$ ]	5.15	16.39	8.75 $\pm$ 0.27	6.220 <sub>47,190</sub>	< 0.0001	1.130 <sub>9,19</sub>	0.29
$E$ [ $\text{mmol m}^{-2} \text{s}^{-1}$ ]	2.19	3.51	2.68 $\pm$ 0.035	2.279 <sub>54,287</sub>	< 0.0001	0.028 <sub>1,287</sub>	0.87
Canopy $E$ [ $\text{mmol s}^{-1}$ ]	0.500	1.480	0.810 $\pm$ 0.026	6.804 <sub>47,190</sub>	< 0.0001	1.408 <sub>10,41</sub>	0.19
$g_s$ [ $\text{mol m}^{-2} \text{s}^{-1}$ ]	0.199	0.356	0.253 $\pm$ 0.004	2.300 <sub>54,287</sub>	< 0.0001	4.459 <sub>1,287</sub>	0.04*
Canopy $g_s$ [ $\text{mol s}^{-1}$ ]	0.047	0.150	0.072 $\pm$ 0.002	7.496 <sub>47,190</sub>	< 0.0001	0.060 <sub>10,90</sub>	0.95
$C_i/C_a$	0.39	0.55	0.47 $\pm$ 0.01	2.398 <sub>54,286</sub>	< 0.0001	-2.803 <sub>27,29</sub>	0.01†
$C_i$	157.9	219.9	190.7 $\pm$ 2.2	2.479 <sub>54,289</sub>	< 0.0001	-2.539 <sub>27,78</sub>	0.02‡
Leaf area [ $\text{m}^2$ ]	0.18	0.50	0.28 $\pm$ 0.01	5.105 <sub>47,190</sub>	< 0.0001	1.244 <sub>9,14</sub>	0.24
Fresh cane mass [g]	678.7	2,051.9	1,230.5 $\pm$ 35.2	4.998 <sub>47,192</sub>	< 0.0001	0.466 <sub>7,65</sub>	0.65
Dry cane mass [g]	190.2	574.6	343.1 $\pm$ 10.2	4.373 <sub>47,192</sub>	< 0.0001	0.774 <sub>7,66</sub>	0.46
Fresh leaf mass [g]	48.9	156.4	88.6 $\pm$ 2.8	7.142 <sub>47,192</sub>	< 0.0001	0.901 <sub>9,33</sub>	0.39
Dry leaf mass [g]	13.8	51.0	29.2 $\pm$ 0.9	5.285 <sub>47,189</sub>	< 0.0001	1.089 <sub>9,52</sub>	0.31
Cane height [cm]	186.2	341.9	269.9 $\pm$ 3.9	12.43 <sub>47,189</sub>	< 0.0001	1.350 <sub>7,14</sub>	0.22
Cane diameter [cm]	2.01	2.93	2.51 $\pm$ 0.03	4.080 <sub>47,192</sub>	< 0.0001	-0.680 <sub>7,08</sub>	0.52
Base cane diameter [cm]	2.31	3.20	2.76 $\pm$ 0.03	2.876 <sub>47,192</sub>	< 0.0001	-1.041 <sub>7,87</sub>	0.33
Mid cane diameter [cm]	1.98	2.80	2.40 $\pm$ 0.03	5.855 <sub>47,192</sub>	< 0.0001	-0.384 <sub>6,84</sub>	0.71
Cane volume [L]	1.005	2.265	1.48 $\pm$ 0.04	4.118 <sub>47,189</sub>	< 0.0001	0.223 <sub>7,98</sub>	0.83
Number of nodes	13.8	23.8	19.0 $\pm$ 0.3	5.013 <sub>47,192</sub>	< 0.0001	-0.009 <sub>11,31</sub>	0.99
Number of green leaves	6.0	10.8	8.3 $\pm$ 0.2	5.901 <sub>47,192</sub>	< 0.0001	-0.935 <sub>7,70</sub>	0.38




Fig. 1. Rank ordered plots of leaf gas-exchange traits. Genotypes (x-axis) are ranked by ascending order for each trait separately. Cumulative  $\text{CO}_2$  assimilation and transpiration represent total  $\text{CO}_2$  assimilation and transpiration that occurred during the time to reach maximum photosynthetic rate (TRMPR). Blue represents mid-stage genotypes currently in the USDA-ARS breeding program in Houma, LA, USA, and red represents commercially released cultivars.  $E$  – transpiration rate;  $g_s$  – stomatal conductance;  $P_N$  – net photosynthetic rate.

photosynthetic capacity. The top commercial cultivar by acreage planted in Louisiana, L 01-299, was in the quadrant of low  $P_N$  and  $g_s$  along with HoCP 20-532, HoCP 17-701, and HoCP 20-534. Commercial cultivars had slightly

higher  $g_s$ ,  $C_i$ , and  $C_i/C_a$ , indicated similar photosynthetic enzyme activities and mesophyll conductance. This relationship strengthened considerably when scaled to the entire canopy (Fig. 2B;  $R^2 = 0.82$ ,  $P < 0.0001$ ).




Fig. 2. Net photosynthetic rate ( $P_N$ ) at the leaf level (A) and canopy level (B) vs. stomatal conductance ( $g_s$ ) at the leaf and canopy levels, respectively of 55 sugarcane genotypes. Blue represents mid-stage genotypes currently in the USDA-ARS breeding program in Houma, LA, USA, and red represents commercially released cultivars. Horizontal and vertical lines represent the mean  $g_s$  and  $P_N$ , respectively. Quadrant I through IV represent combinations of high and low  $P_N$  and  $g_s$ . Canopy level gas exchange was calculated as gas exchange for the entire canopy leaf area, assuming uniform gas exchange over the entire canopy.

Stomatal propensity to remain open varied substantially across genotypes as expressed by genotypic variation in TRMPR, showing a significant genotype effect. TRMPR ranged from 28.7 to 111 min, showing a nearly fourfold difference across genotypes. The trait was somewhat consistent among replicates of the same genotype with standard errors between 1.5 to 34% (mean 10.9%) of the TRMPR value. TRMPR was not significantly correlated to  $E$  or  $g_s$ , but  $P_N$  and TRMPR formed a significant positive correlation, meaning that genotypes with large maximum  $P_N$  also reached maximum  $P_N$  more quickly ( $P < 0.05$ ). Further, the regression between maximum  $P_N$  and the cumulative  $\text{CO}_2$  assimilation [ $\mu\text{mol m}^{-2}$ ] that took place while the leaf was reaching maximum  $P_N$  was significant with an  $R^2$  of 0.79 (Fig. 3A;  $P < 0.0001$ ). When the cumulative  $\text{CO}_2$  assimilation was regressed on TRMPR, the regression was significant ( $R^2 = 0.36$ ), meaning that TRMPR accounted for 21–36% of the variation in the cumulative  $\text{CO}_2$  assimilation. Greater cumulative photosynthesis took place when the leaf reached maximal  $P_N$  more rapidly (Fig. 3B). In contrast,  $E$  and  $g_s$  regressed on the cumulative  $E$  and  $g_s$ , respectively, resulting in  $R^2$  of 0.89 and 0.90, respectively, while TRMPR was not significantly correlated to cumulative  $E$  and  $g_s$ , showing that TRMPR did not affect that total water loss from the leaves (Fig. 3C,D). However, TRMPR was significantly negatively correlated with the initial  $g_s$  when the leaf was first placed in the gas-exchange cuvette ( $R^2 = 0.21$ ,  $P = 0.0006$ ), where short TRMPR was associated with high initial  $g_s$ . This relationship increased in strength over the first 10 min where the correlation between  $g_s$  and TRMPR reached a plateau at  $R^2$  of 0.3–0.35 ( $P < 0.0001$ ). Genotypes, HoCP 20-556, HoCP 18-829, Ho 18-878, L 17-428, and HoCP 14-885 had the shortest TRMPR, while HoCP 20-541, Ho 13-739, HoCP 20-532, HoCP 20-535, and HoCP 20-529 had the longest TRMPR (Table 1S). The sugarcane genotypes that ranked high in

$P_N$  and TRMPR were L 17-428, Ho 17-738, HoCP 20-505, HoCP 14-885, and L 12-201, while HoCP 20-532, HoCP 20-529, HoCP 20-501, HoCP 20-535, and HoCP 20-527 were the lowest (Table 1S).

**Whole-plant traits:** Whole-plant traits exhibited more variation across genotypes than gas-exchange traits in that maximum values were 1.46 to 3.70 times that of the minimum values. This was particularly true with mass traits and leaf area (3.2 $\times$ ) with less variability in cane height and diameter and number of nodes (1.67 $\times$ ). Like gas-exchange traits, the traits were consistent across replicates of each genotype, where standard errors were 2–8% of the measurement values. Mass traits had greater variation with SE representing 7–8% of the measurement, while cane diameter, height, and number of nodes were only 2–4% of the measurement. All whole-plant traits were significantly different across genotypes, showing a significant genotype effect (Fig. 4). The commercial cultivars and unreleased genotypes in the breeding program did not differ in any whole-plant trait ( $P > 0.05$ ; Table 1, Fig. 4). A positive relationship was found between stalk and canopy traits, where stalks with greater leaf area ( $r = 0.61$ ) and leaf mass ( $r = 0.65$ ) tended to be heavier (Fig. 5;  $P < 0.0001$ ). No relationship was found between gas-exchange traits and whole-plant traits.

The sugarcane genotypes that ranked highest in every whole-plant trait were HoCP 20-541, HoCP 20-523, HoCP 20-538, HoCP 14-885, and L 17-428. HoCP 18-829, HoCP 20-525, L 14-267, HoCP 20-502, Ho 18-878, and HoCP 20-548 consistently ranked low. Overall, in both gas exchange and whole-plant traits, the highest ranked genotypes were L 17-428, L 12-201, HoCP 20-523, HoCP 14-885, and HoCP 20-556, while the lowest ranked genotypes were HoCP 18-829, HoCP 20-502, HoCP 17-701, L 01-299, and HoCP 20-878. The rankings for gas exchange and whole-plant traits were not correlated, even




Fig. 3. Relationship of cumulative photosynthesis during the time to reach maximum photosynthetic rate (TRMPR) with  $P_N$  (A) and TRMPR (B) and cumulative  $E$  and  $g_s$  during TRMPR vs.  $E$  (C) and  $g_s$  (D) at TRMPR, respectively. Blue represents mid-stage genotypes currently in the USDA-ARS breeding program in Houma, LA, USA, and red represents commercially released cultivars. TRMPR and either  $E$  or  $g_s$  at TRMPR were not significantly correlated.  $E$  – transpiration rate;  $g_s$  – stomatal conductance;  $P_N$  – net photosynthetic rate.

though several genotypes were found at the top of each list. Multiple genotypes had low stalk mass but large canopies both in leaf area and mass such as HoCP 20-510 and HoCP 20-563 or large cane mass but small canopies such as HoCP 20-504 and HoCP 20-512 (Table 2S, *supplement*).

## Discussion

Gas-exchange traits exhibited relatively large variation even in this highly selected group of commercial cultivars and mid-stage genotypes in the breeding program. Breeding for early sucrose accumulation, sugar and fresh aboveground biomass yield per hectare, and disease resistance has substantially increased sugar production over the previous 120 years (Hale *et al.* 2022). In sugarcane, similar to many other crops, substantial yield improvements have been made by selecting for increased carbon allocation to harvestable biomass and increased harvest index (Long *et al.* 2006, Zhu *et al.* 2010). The fact that other traits such as gas-exchange traits are not directly selected and still exhibit significant variation across genotypes indicates that multiple different phenotypic and genetic scenarios can result in a relatively good sugarcane cultivar. A lack of difference between commercial and unreleased mid-stage

genotypes further indicates that likely photosynthetic traits are not being selected indirectly either. Therefore, selecting genotypes with high  $P_N$  or short TRMPR that increase carbon inputs along with the traditional breeding targets that select for increased carbon allocation to sugar production can result in greater potential yields than selecting for increased carbon allocation to sugar alone. Genetic manipulation and redesigning crop plants are often considered necessary avenues to increase potential yields through improvements in plant physiology (Zhu *et al.* 2010, Long *et al.* 2015, Ort *et al.* 2015), but substantial variability in gas-exchange traits in sugarcane indicates that traditional selection can still increase genetic gain. Even slight improvements in photosynthesis of 2–3% can produce noticeable yield increases by harvest time (Lefebvre *et al.* 2005, Lawson and Blatt 2014).

Variation in the interaction between gas-exchange traits,  $P_N$  and  $g_s$ , suggests that sugarcane exhibits variation in both photosynthetic capacity and  $\text{CO}_2$  substrate availability. Other studies have also recorded variation in photosynthetic capacity and  $g_s$  in sugarcane, indicating an opportunity to select for improved photosynthesis (Inman-Bamber *et al.* 2011, Hoffman *et al.* 2018, Zhao *et al.* 2019, Singels *et al.* 2021). Differences in




Fig. 4. Rank ordered plots of whole-plant traits. Genotypes (x-axis) are ranked by ascending order for each trait separately. Cane diameter is the mean of diameters measured at base and mid distance on the cane. Fresh and dry cane mass represent the mass of the stalk only without leaves. *Blue* represents mid-stage genotypes currently in the USDA-ARS breeding program in Houma, LA, USA, and *red* represents commercially released cultivars.




Fig. 5. Cane mass vs. canopy leaf area (A) and mass (B). *Blue* represents mid-stage genotypes currently in the USDA-ARS breeding program in Houma, LA, USA, and *red* represents commercially released cultivars.

photosynthetic capacity were identified as genotypes that had high  $P_N$  and low  $C_i/C_a$  compared to  $g_s$ , which could indicate a physiological difference such as modified bundle-sheath leakiness or higher mesophyll conductance or PEPC and Rubisco activities (Kromdijk *et al.* 2014, Li *et al.* 2017, Lawson *et al.* 2018).  $C_i$  and  $C_i/C_a$  are heritable in sugarcane and have been considered a proxy for increased water-use efficiency ( $P_N/g_s$ ) because they indicate increased photosynthetic capacity (and, therefore,  $P_N$ ) relative to  $g_s$ , but these proxies may provide a heritable screen of photosynthetic capacity, especially when coupled with increased  $P_N$  (Ghannoum 2016, Jackson *et al.* 2016). Increased substrate availability is defined as having increased  $CO_2$  diffusion into the stomata (high  $g_s$ ), resulting in higher  $P_N$  (Li *et al.* 2017). For example,

Ho 17-738 appears to have high photosynthetic capacity, while L 17-428 had very high  $g_s$  to increase  $CO_2$  substrate availability to fuel the highest  $P_N$  measured in the study and relatively low  $C_i/C_a$ , potentially having both increased photosynthetic capacity and substrate availability. The current top cultivar by total acreage planted in Louisiana, L 01-299, has multiple ideal traits such as ratooning ability that have maintained its importance in the sugarcane industry, yet it had rather low  $P_N$  and  $g_s$  (Gravois *et al.* 2011). Incorporating improved photosynthetic traits in an excellent cultivar could lead to greater carbon inputs that can be converted to sugar production. Increasing substrate availability by greater  $g_s$  has the additional consequence of increased water loss (Lawson and Violet-Chabrand 2019), but despite that

higher substrate availability is an important trait because sugarcane seldom lack sufficient water in water-abundant south Louisiana (Ellsworth and White Jr. 2022).

Increased CO<sub>2</sub> assimilation during transitory periods such as fluctuating light can improve total photosynthetic output even if accompanied with increased water loss. Theoretically, stomata are only open when conditions are right for photosynthesis, attempting to adjust stomatal activity with photosynthetic demands to reduce water loss and maximize water-use efficiency (Brodribb *et al.* 2009, McAusland *et al.* 2016, Lawson and Vialet-Chabrand 2019). Considering that photosynthesis during disturbance and fluctuating light conditions account for at least 35% of all fixed carbon, closing stomata under fluctuating light and other disturbances is a major limiting factor for photosynthesis and increasing photosynthesis during transitory periods through increasing the propensity of stomata to remain open can increase photosynthetic output beyond the effect of greater photosynthetic capacity alone (Leakey *et al.* 2002, Lawson *et al.* 2018, Condon 2020, Yamori *et al.* 2020, Wang *et al.* 2021). Genotypes with the shortest TRMPR values had the highest initial  $g_s$ , so TRMPR trait represents the propensity of stomata to remain partially open during disturbance and to quickly reopen completely after disturbance. TRMPR could lead to increased transpiration, but it does not appear that TRMPR affects cumulative  $E$  as much as it affects cumulative  $P_N$  during transitory periods. Considering that fluctuating light is common for much of the leaf area in the canopy, increasing TRMPR could improve photosynthesis outputs in sugarcane in Louisiana where water abundance and high humidity do not necessitate water conservation.

Combining the selection of leaf- and whole plant-level traits such as photosynthesis and canopy traits can create a synergistic effect. For example, the photosynthetic output of the whole canopy across genotypes was strongly influenced by canopy size, leading to a stronger relationship between  $P_N$  and  $E$ , where some large-canopied varieties had high canopy  $P_N$  or  $E$ , even though leaf-level  $P_N$  or  $E$  were low. This potentially shows variation in the hydraulic constraints on gas exchange and the need to understand photosynthetic traits across scales (Meinzer and Grantz 1990, Sperry 2000). The fact that increased canopy leaf area and mass was correlated with increased stalk mass and volume indicates that coupling high photosynthesis with large canopy leaf area could greatly increase aboveground biomass and sugar production. Interestingly, this relationship between leaf area and stalk size appears to contradict the concept of an ideal canopy size where optimal canopy leaf area is less than maximal canopy size (Srinivasan *et al.* 2017). For example, biomass productivity of *Miscanthus × giganteus* in the Midwest USA exceeded maize (*Zea mays*) because it reached 90% canopy closure about four weeks before maize (Dohleman and Long 2009). Sugarcane grows slowly and takes longer to reach canopy closure, resulting in inefficient light interception for the first part of the growing season (Allison *et al.* 2007, Inman-Bamber 2014, Dias *et al.* 2020). Therefore, an ideal sugarcane canopy would include vigorous initial growth and large size to reach canopy closure quickly and

leaves with high photosynthetic capacity and low TRMPR to increase photosynthetic output throughout the growing season, leading to greater biomass and sugar yields (Zhu *et al.* 2010).

No relationship was found between gas exchange and whole-plant traits, but that does not indicate that leaf-level traits do not influence whole-plant traits; rather it shows the complexity of factors that hinder the scaling of traits. Because CO<sub>2</sub> assimilation and water loss from the stomata are essentially the only source of C gain and water loss, there is an implicit relationship between leaf-level gas exchange and whole-plant growth and water loss (Farquhar *et al.* 1989, Singels *et al.* 2021). Nonetheless, this relationship can be obscured because of fixed carbon losses such as non-leaf respiration and differences in carbon use such as allocation to roots. Therefore, sink strength often is stronger than source strength (photosynthesis), so that sugar transport and accumulation limit yield more than photosynthesis (McCormick *et al.* 2006, 2008, 2009; Evans 2013, Singels *et al.* 2021). Additionally, single point measurements of gas exchange do not necessarily represent mean photosynthetic rate and water loss for the whole canopy, and comparisons across genotypes conflate gas exchange differences with genotypic whole plant and leaf variation (Long *et al.* 2006). However, a clear correlation between  $P_N$  and  $g_s$  with yield was observed in sugarcane in Florida, USA, and in other C<sub>4</sub> Panicoid grass species (Zhao *et al.* 2015, 2019; Ellsworth *et al.* 2017). As a stronger connection between leaf-level photosynthetic traits and whole-plant traits becomes better understood, we will be better able to identify suitable traits, leading to better phenotyping and increased genetic gain.

**Conclusion:** Improving photosynthetic traits is one of the underutilized areas in plant breeding, and they possess great potential to increase carbon inputs. Gas exchange is a powerful tool to measure multiple traits related to CO<sub>2</sub> assimilation and water loss. Nonetheless, more work needs to be done to make better connections between leaf- and whole plant-level traits, so that explicit relationships can be defined and used as a basis for selection. In humid, water-abundant agroecosystems, stomatal propensity to remain open (short TRMPR) has potential to improve photosynthesis through increasing CO<sub>2</sub> availability in frequently occurring transitory periods. Coupled with greater photosynthetic capacity, this stomatal propensity to remain open can substantially increase photosynthesis in sugarcane. High-throughput phenotyping is problematic in that these measurements are time-consuming, so these measurements may be most useful in a more advanced stage of selection in the breeding program.

## References

- Allison J.C.S., Pammenter N.W., Haslam R.J.: Why does sugarcane (*Saccharum* sp. hybrid) grow slowly? – S. Afr. J. Bot. **73**: 546-551, 2007.
- Araus J.L., Kefauver S.C., Zaman-Allah M. *et al.*: Translating high-throughput phenotyping into genetic gain. – Trends Plant Sci. **23**: 451-466, 2018.
- Araus J.L., Li J., Parry M.A.J., Wang J.: Phenotyping and other

- breeding approaches for a New Green Revolution. – *J. Integr. Plant Biol.* **56**: 422-424, 2014.
- ASCL: The Louisiana Sugar Industry – American Sugar Cane League, 2023. Available at: <https://www.amscl.org/about-the-league/>.
- Brodribb T.J., McAdam S.A.M., Jordan G.J., Feild T.S.: Evolution of stomatal responsiveness to CO<sub>2</sub> and optimization of water-use efficiency among land plants. – *New Phytol.* **183**: 839-847, 2009.
- Condon A.G.: Drying times: plant traits to improve crop water use efficiency and yield. – *J. Exp. Bot.* **71**: 2239-2252, 2020.
- Dias H.B., Inman-Bamber G., Everingham Y. *et al.*: Traits for canopy development and light interception by twenty-seven Brazilian sugarcane varieties. – *Field Crop. Res.* **249**: 107716, 2020.
- Dohleman F.G., Long S.P.: More productive than maize in the Midwest: how does *Miscanthus* do it? – *Plant Physiol.* **150**: 2104-2115, 2009.
- Ellsworth P.Z., Ellsworth P.V., Cousins A.B.: Relationship of leaf oxygen and carbon isotopic composition with transpiration efficiency in the C<sub>4</sub> grasses *Setaria viridis* and *Setaria italica*. – *J. Exp. Bot.* **68**: 3513-3528, 2017.
- Ellsworth P.Z., Feldman M.J., Baxter I., Cousins A.B.: A genetic link between whole-plant water use efficiency and leaf carbon isotope composition in the C<sub>4</sub> grass *Setaria*. – *Plant J.* **102**: 1234-1248, 2020.
- Ellsworth P.Z., White Jr. P.M.: Row spacing and the use of plant-available water in sugarcane cultivation in water-abundant Louisiana. – *Agronomy* **12**: 1586, 2022.
- Evans J.R.: Improving photosynthesis. – *Plant Physiol.* **162**: 1780-1793, 2013.
- Farquhar G.D., Hubick K.T., Condon A.G., Richards R.A.: Carbon isotope fractionation and plant water-use efficiency. – In: Rundel P.W., Ehleringer J.R., Nagy K.A. (ed.): *Stable Isotopes in Ecological Research*. Pp. 21-40. Springer, New York 1989.
- Feldman M.J., Ellsworth P.Z., Fahlgren N. *et al.*: Components of water use efficiency have unique genetic signatures in the model C<sub>4</sub> grass *Setaria*. – *Plant Physiol.* **178**: 699-715, 2018.
- Ghannoum O.: How can we breed for more water use-efficient sugarcane? – *J. Exp. Bot.* **67**: 557-559, 2016.
- Gravois K.A., Bischoff K.P., Pontif M.J. *et al.*: Registration of ‘L 01–299’ sugarcane. – *J. Plant Regist.* **5**: 191-195, 2011.
- Hale A.L., Todd J.R., Gravois K.A. *et al.*: Sugarcane breeding programs in the USA. – *Sugar Tech.* **24**: 97-111, 2022.
- Hoffman N., Singels A., Patton A., Ramburan S.: Predicting genotypic differences in irrigated sugarcane yield using the Canegro model and independent trait parameter estimates. – *Eur. J. Agron.* **96**: 13-21, 2018.
- Inman-Bamber G.: Sugarcane yields and yield-limiting processes. – In: Moore P.H., Botha F.C. (ed.): *Sugarcane: Physiology, Biochemistry, and Functional Biology*. Pp. 579-600. John Wiley & Sons, Ames 2014.
- Inman-Bamber N.G., Jackson P.A., Hewitt M.: Sucrose accumulation in sugarcane stalks does not limit photosynthesis and biomass production. – *Crop Pasture Sci.* **62**: 848-858, 2011.
- ISO: Sugar Sector – International Sugar Organization, 2023. Available at: <https://www.isosugar.org/sugarsector/sugar>.
- Jackson P., Basnayake J., Inman-Bamber G. *et al.*: Genetic variation in transpiration efficiency and relationships between whole plant and leaf gas exchange measurements in *Saccharum* spp. and related germplasm. – *J. Exp. Bot.* **67**: 861-871, 2016.
- Kromdijk J., Ubierna N., Cousins A.B., Griffiths H.: Bundle sheath leakiness in C<sub>4</sub> photosynthesis: a careful balancing act between CO<sub>2</sub> concentration and assimilation. – *J. Exp. Bot.* **65**: 3443-3457, 2014.
- Lawson T., Blatt M.R.: Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. – *Plant Physiol.* **164**: 1556-1570, 2014.
- Lawson T., Terashima I., Fujita T., Wang Y.: Coordination between photosynthesis and stomatal behavior. – In: Adams W.W., Terashima I. (ed.): *The Leaf: A Platform for Performing Photosynthesis*. Pp. 141-161. Springer, Cham 2018.
- Lawson T., Viallet-Chabrand S.: Speedy stomata, photosynthesis and plant water use efficiency. – *New Phytol.* **221**: 93-98, 2019.
- Leakey A.D.B., Ferguson J.N., Pignou C.P. *et al.*: Water use efficiency as a constraint and target for improving the resilience and productivity of C<sub>3</sub> and C<sub>4</sub> crops. – *Annu. Rev. Plant Biol.* **70**: 781-808, 2019.
- Leakey A.D.B., Press M.C., Scholes J.D., Watling J.R.: Relative enhancement of photosynthesis and growth at elevated CO<sub>2</sub> is greater under sunflecks than uniform irradiance in a tropical rain forest tree seedling. – *Plant Cell Environ.* **25**: 1701-1714, 2002.
- Lefebvre S., Lawson T., Fryer M. *et al.*: Increased sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis and growth from an early stage in development. – *Plant Physiol.* **138**: 451-460, 2005.
- Li C., Jackson P., Lu X. *et al.*: Genotypic variation in transpiration efficiency due to differences in photosynthetic capacity among sugarcane-related clones. – *J. Exp. Bot.* **68**: 2377-2385, 2017.
- Long S.P., Bernacchi C.J.: Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. – *J. Exp. Bot.* **54**: 2393-2401, 2003.
- Long S.P., Marshall-Colon A., Zhu X.-G.: Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. – *Cell* **161**: 56-66, 2015.
- Long S.P., Zhu X.-G., Naidu S.L., Ort D.R.: Can improvement in photosynthesis increase crop yields? – *Plant Cell Environ.* **29**: 315-330, 2006.
- McAusland L., Viallet-Chabrand S., Davey P. *et al.*: Effects of kinetics of light-induced stomatal responses on photosynthesis and water-use efficiency. – *New Phytol.* **211**: 1209-1220, 2016.
- McCormick A.J., Cramer M.D., Watt D.A.: Sink strength regulates photosynthesis in sugarcane. – *New Phytol.* **171**: 759-770, 2006.
- McCormick A.J., Cramer M.D., Watt D.A.: Regulation of photosynthesis by sugars in sugarcane leaves. – *J. Plant Physiol.* **165**: 1817-1829, 2008.
- McCormick A.J., Watt D.A., Cramer M.D.: Supply and demand: sink regulation of sugar accumulation in sugarcane. – *J. Exp. Bot.* **60**: 357-364, 2009.
- Medrano H., Tomás M., Martorell S. *et al.*: From leaf to whole-plant water use efficiency (WUE) in complex canopies: Limitations of leaf WUE as a selection target. – *Crop J.* **3**: 220-228, 2015.
- Meinzer F.C., Grantz D.A.: Stomatal and hydraulic conductance in growing sugarcane: stomatal adjustment to water transport capacity. – *Plant Cell Environ.* **13**: 383-388, 1990.
- Ort D.R., Merchant S.S., Alric J. *et al.*: Redesigning photosynthesis to sustainably meet global food and bioenergy demand. – *PNAS* **112**: 8529-8536, 2015.
- R Core Team: *R: A Language and Environment for Statistical Computing*. R Version 4.3.1, R Foundation for Statistical Computing, 2021.
- Singels A., Jackson P., Inman-Bamber G.: Sugarcane. – In: Sadras

- V.O., Calderini D.F. (ed.): Crop Physiology: Case Histories for Major Crops. Pp. 674-713. Academic Press, London 2021.
- Sperry J.S.: Hydraulic constraints on plant gas exchange. – *Agr. Forest Meteorol.* **104**: 13-23, 2000.
- Srinivasan V., Kumar P., Long S.P.: Decreasing, not increasing, leaf area will raise crop yields under global atmospheric change. – *Glob. Change Biol.* **23**: 1626-1635, 2017.
- von Caemmerer S.: Biochemical Models of Leaf Photosynthesis. Pp. 165. Csiro Publishing, Collingwood 2000.
- von Caemmerer S., Farquhar G.D., Berry J.A.: Biochemical model of C<sub>3</sub> photosynthesis. – In: Laisk A., Nedbal L., Govindjee (ed.): *Photosynthesis in silico: Understanding Complexity from Molecules to Ecosystems. Advances in Photosynthesis and Respiration.* Vol. 29. Pp. 209-230. Springer, Dordrecht 2009.
- von Caemmerer S., Furbank R.T.: The C<sub>4</sub> pathway: an efficient CO<sub>2</sub> pump. – *Photosynth. Res.* **77**: 191-207, 2003.
- Wang Y., Chan K.X., Long S.P.: Towards a dynamic photosynthesis model to guide yield improvement in C<sub>4</sub> crops. – *Plant J.* **107**: 343-359, 2021.
- Yamori W., Kusumi K., Iba K., Terashima I.: Increased stomatal conductance induces rapid changes to photosynthetic rate in response to naturally fluctuating light conditions in rice. – *Plant Cell Environ.* **43**: 1230-1240, 2020.
- Zhao D., Glaz B., Irej M.S., Hu C.-J.: Sugarcane genotype variation in leaf photosynthesis properties and yield as affected by mill mud application. – *Agron. J.* **107**: 506-514, 2015.
- Zhao D., Irej M., LaBorde C., Hu C.-J.: Physiological and yield characteristics of 18 sugarcane genotypes grown on a sand soil. – *Crop Sci.* **59**: 2741-2751, 2019.
- Zhu X.-G., Long S.P., Ort D.R.: Improving photosynthetic efficiency for greater yield. – *Annu. Rev. Plant Biol.* **61**: 235-261, 2010.

© The authors. This is an open access article distributed under the terms of the Creative Commons BY-NC-ND Licence.